



American Embassy School
New Delhi

Chandragupta Marg
Chanakyapuri, New Delhi 110021 INDIA
[p]+91 11 2688 8854 [f] +91 11 2687 3320
[w] www.aes.ac.in

**HEALTH
FORM 2
IMMUNIZATION**

Immunization History

Student Name _____ Age _____ Date _____

Provide All Dates -month/day/year. Gray boxes are not required.

Required	#1	#2	#3	#4	#5	TD (over age 7)
Diphtheria, Tetanus, Pertussis -DTP, DTaP, TDaP or DT						
Poliomyelitis						
Hib (Haemophilus Influenza Type b) - Required for all Pre-K and Kindergarten students						
MMR (Measles, Mumps, Rubella)						
Measles (if given separately)						
Hepatitis B						
Recommended	Please fill in the dates, if any of these were given.					
Hepatitis A						
Meningococcal						
Varicella (chickenpox)						
Typhoid						
Rabies						
Japanese Encephalitis						
HPV (Girls and Boys older than 13 yrs.)						
PCV (Pneumococcal)						
Rotavirus						

Has your child contracted measles or varicella (chicken pox)? If yes, which year? _____

NOTE: It is the responsibility of the parent/guardian to notify the Health Office of any changes to the information given.

Signature of Parent _____ Date _____

AES IMMUNIZATION AND TB SCREENING REQUIREMENTS

IMMUNIZATIONS REQUIRED FOR ATTENDANCE AT AES:

DPT, DtaP, or Td or Tdap (Diphtheria and tetanus, with or without pertussis)—A minimum of 4 doses, with at least one dose after the 4th birthday. Middle and High School students whose last immunization containing tetanus was 10 or more years ago must receive a Td (adult tetanus-diphtheria) or Tdap (adolescent tetanus, diphtheria and pertussis booster) prior to attending school sponsored travel such as WOW, Minicourse, ASIAC/MESAC or field trips.

POLIO—4 doses of oral or injectable polio. If one dose has been administered after the 4th birthday, then the only 3 doses are required.

Hib (Haemophilus influenza type B)—Required for children below 5 years of age are (PK and KG only)

MEASLES, MUMPS, RUBELLA—PK and KG requires one dose of MMR. Grade 1 and up require 2 two doses of MMR or 1 Measles and 1 MMR.

HEPATITIS B—Series of 3 required from all the students.

All dates from birth must be provided. Students who do not have immunization records may be requested to have a repeat immunization of one of each of the above, to ensure at least partial immunity.

TUBERCULOSIS (TB) SCREENING: Students are required to have tuberculosis screening within 3 months prior to admission to AES and every two years thereafter.

No BCG (TB) vaccination

Tuberculosis screening should be done on admission and every 2 years by a licensed practitioner. This is completed by-

- 1) a PPD Mantoux test **or**
- 2) a chest x-ray (CXR) **or**
- 3) a blood test-Quantiferon

If a student has history of a positive PPD Mantoux test and has completed the treatment course they would require a chest X-ray for clearance.

With BCG (TB) vaccination

Tuberculosis screening should be done on admission and every 2 years by a licensed practitioner. This is completed by-
An examination of the chest to confirm no active symptoms of TB

If a student has had their BCG vaccination within the last 5 years, they do not require a TB screening; please make sure the Health Office has the BCG date recorded. BCG (TB) vaccinations are not required for AES.

ADDITIONAL IMMUNIZATIONS RECOMMENDED:

AES does not require these for admission, but they are highly recommended for the health of your children.

VARICELLA- (chickenpox)—2 doses for all ages

TYPHOID—Typhoid is spread through contaminated food or water. One shot is good for 2-3 years or 4 tablets orally, good for 5 years. Repeat at recommended time. The oral vaccine is not available in India.

HEPATITIS A- Hepatitis A is spread through contaminated food or water. 2 doses administered 6 months apart.

RABIES Pre Exposure—3 doses in one month (Days 1, 7, 21 or 28). Rabies is very prevalent all over India. The CDC recommends people who are taking long trips or moving to India to have rabies vaccinations. They emphasize the importance of vaccinating children, because they tend to play with animals, might not report bites, and are more likely to have animal bites on their head and neck.

JAPANESE ENCEPHALITIS—2 doses in 28 days apart. Serious illness carried by mosquitoes. Not usually present in urban areas. Recommended for those students likely to visit rural areas with rice and livestock production, particularly eastern coastal states, and parts of Nepal.

MENINGOCOCCAL- 1 dose of this vaccine for children ages 11-19 years of age. There are different strains of this bacterial meningitis that can cause serious illness or death in different countries therefore consult your doctor. A booster may be required every 3 years depending on vaccine.

(HPV) Human Papillomavirus- HPV vaccine is important because it protects against cancers caused by HPV infection. All children who are 11 or 12 years old should get two shots of HPV vaccine six to twelve months apart. Adolescents who receive their two shots less than five months apart will require a third dose of HPV vaccine.